

Common Carp (*Cyprinus carpio*)

Ecological Risk Screening Summary

Web Version—09/10/2014




Photo: © www.sportviszone.nl from Frankis et al. (2014).

1 Native Range, and Status in the United States

Native Range

From Kottelat and Freyhof (2007):

“Europe to Asia: Black, Caspian and Aral Sea basins. Introduced throughout the world. Wild stocks are only present naturally in rivers draining to the Black, Caspian and Aral Sea. A reophilic wild population in the Danube is assumed to be the origin of the European species; this population is now under threat (Kottelat 1997).”

Status in the United States

From Nico et al. (2014):

“Recorded from all states except Alaska. In their summary table, Bailey and Smith (1981) indicated that *Cyprinus carpio* is widely distributed in the Great Lakes basin.”

“Records of introductions are available for the following states: Alabama (Smiley 1886, Baird 1887, McDonald 1886, 1893, Spencer et al. 1964, Smith-Vaniz 1968, Dahlberg and Scott 1971a, Swift et al. 1977, Mettee et al. 1987, 1996, Boschung 1992, Rasmussen 1998); Arizona (Rule

1885, Taggart 1885, Evermann and Rutter 1895, Gilbert and Scofield 1898, Miller and Lowe 1967, Minckley 1973, Lee et al. 1980 et seq., Tilmant 1999, USFWS 2005, Illinois Natural History Survey Fish Collections); Arkansas (Baird 1887, McDonald 1887, 1893, Courtenay 1970, Lee et al. 1980 et seq., Robison and Buchanan 1988, Rasmussen 1998, USFWS 2005, Mississippi Museum of Natural Science 2004, Illinois Natural History Survey Fish Collections); California (Smiley 1886, Smith 1896, Shebley 1917, Lampman 1946, La Rivers 1962, Moyle et al. 1974, Moyle 1976, Lee et al. 1980 et seq., Moyle and Daniels 1982, Smith 1982, Tilmant 1999, Sommer et al. 2001, Moyle 2002, USFWS 2005, Matern [et al.] 2002, Illinois Natural History Survey Fish Collections); Colorado (Smiley 1886, Baird 1887, McDonald 1886, Ellis 1974, Wiltzius 1981, Woodling 1985, Zuckerman and Behnke 1986, Rasmussen 1998, Tilmant 1999, Illinois Natural History Survey 2004); Connecticut (Smiley 1886, Ravenel 1896, Webster 1941, Lee et al. 1980 et seq., Whitworth 1996); Delaware (Smiley 1886, Baird 1887, McDonald 1886, Lee et al. 1976, Raasch and Altemus 1991, USFWS 2005); District of Columbia (Tilmant 1999); Florida (Baird 1887, McDonald 1886, Anonymous 1892, Courtenay et al. 1974, Swift et al. 1977, Shafland 1996, Anonymous 2001, Nico 2005); Georgia (Smiley 1886, Baird 1887, McDonald 1886, 1893, Anonymous 1892, Worth 1895, Ravenel 1896, 1898, Hildebrand 1923, Dahlberg and Scott 1971a, 1971b, Burkhead et al. 1997, Walters 1997); Hawaii (Cobb 1902, Jordan and Evermann 1902, 1905, Brock 1960, Maciolek 1984, Devick 1991, Tilmant 1999, Mundy 2005); Idaho (Smith 1896, Lampman 1946, Linder 1963, Simpson and Wallace 1978, Wydoski and Whitney 1979, Lee et al. 1980 et seq., Idaho Fish and Game 1990, Sigler and Sigler 1996, USFWS 2005, Amercian Fisheries Society 2001); Illinois (Smiley 1886, Baird 1887, McDonald 1886, Hay 1894, Sweeney 1902, Smith 1979, Lee et al. 1980 et seq., Emery 1985, Laird and Page 1996, Rasmussen 1998, Illinois Natural History Survey 2004, USFWS 2005); Indiana (Smiley 1886, Anonymous 1892, Hay 1894, Sweeney 1902, Blatchley 1938, Gerking 1945, Nelson and Gerking 1968, Lee et al. 1980 et seq., Emery 1985, Burr and Page 1986, Tilmant 1999, USFWS 2005); Iowa (Cleary 1956, Bailey and Allum 1962, Lee et al. 1980 et seq., Burr and Page 1986, Harlan et al. 1987, Young et al. 1997, USFWS 2005, Rasmussen, unpublished data); Kansas (Smiley 1886, Anonymous 1892, Ravenel 1896, Dyche 1914, Breukelman 1946, Call 1961, Cross 1967, Lee et al. 1980 et seq., Cross and Collins 1995, Rasmussen 1998, Tilmant 1999); Kentucky (McDonald 1893, Worth 1895, Ravenel 1896, Clay 1975, Lee et al. 1980 et seq., Burr and Page 1986, Burr and Warren 1986, Powers and Ceas 2000); Louisiana (Baird 1887, McDonald 1886, Anonymous 1892, Douglas and Davis 1967, Douglas 1974, Lee et al. 1980 et seq., Piler personal communication.); Maine (Everhart 1976); Maryland (Ferguson 1876, Smiley 1886, McDonald 1886, 1893, Anonymous 1892, Ravenel 1898, Truitt et al. 1929, Schwartz 1963, Lee et al. 1976, 1980 et seq., Tilmant 1999, Starnes et al. 2011); Massachusetts (McDonald 1886, Baird 1887, Lee et al. 1980 et seq., Hartel 1992, Hartel et al. 1996, Bozeman and Charp 2001, USFWS 2005); Michigan (Smiley 1886, McDonald 1893, Hubbs and Cooper 1936, Hubbs and Lagler 1958, Emery 1985, Tilmant 1999, Cudmore-Vokey and Crossman 2000, University of Michigan Museum of Zoology 2004); Minnesota (Baird 1887, McDonald 1886, 1893, Moore and Bream 1965, Eddy and Underhill 1974, Phillips et al. 1982, Emery 1985, Burr and Page 1986, Rasmussen 1998, Tilmant 1999, Myers 2004, Minnesota Sea Grant 2004, USFWS 2005); Mississippi (Smiley 1886, Baird 1887, McDonald 1886, 1893, Cook 1959, Ross and Brenneman 1991, Schramm and Basler 2004, Mississippi Museum of Natural Science 2004); Missouri (Smiley 1886, McDonald 1893, Ravenel 1896, 1898, Pflieger 1971, 1975, 1997, Burr and Page 1986, Young et al. 1997, Rasmussen 1998, USFWS 2005, Mississippi Museum of Natural Science 2004); Montana (Brown 1971, Courtenay 1985, Holton

1990, Young et al. 1997, Tilmant 1999, USFWS 2005); Nebraska (Bailey and Allum 1962, Morris et al. 1974, Lee et al. 1980 et seq., Texas Parks and Wildlife Department. 2001, Nebraska Parks and Wildlife Commission, personal communication); Nevada (Smith 1896, Miller and Alcorn 1946, Miller 1952, Lampman 1946, La Rivers 1962, Bradley and Deacon 1967, Deacon and Williams 1984, Scopettone et al. 1998, Tilmant 1999, Insider Viewpoint 2001, USFWS 2005, Vinyard 2001); New Hampshire (Scarola 1973); New Jersey (Smiley 1886, Nelson 1890, Ravenel 1898, Fowler 1906, 1952, Stiles 1978, USFWS 2005); New Mexico (McDonald 1886, Baird 1887, Koster 1957, Lee et al. 1980 et seq., Sublette et al. 1990, Platania 1991, New Mexico Game and Fish 2000); New York (Smiley 1886, McDonald 1893, Bean 1903, Lee et al. 1980 et seq., Werner 1980, Emery 1985, Smith 1985); North Carolina (Anonymous 1892, Worth 1895, Ravenel 1896, Cahoon 1953, Lee et al. 1980 et seq., Menhinick 1991, USFWS 2005); North Dakota (Owen et al. 1981, Young et al. 1997, Power and Ryckman 1998, USFWS 2005); Ohio (Jordan 1882, Smiley 1886, Baird 1887, McDonald 1886, 1893, Anonymous 1892, Trautman 1981, Emery 1985, Burr and Page 1986, Tilmant 1999, USFWS 2005); Oklahoma (McDonald 1893, Bean 1896, Hall 1956, Miller and Robison 1973, Lee et al. 1980 et seq., Rasmussen 1998); Oregon (Smith 1896, Lampman 1946, Wydoski and Whitney 1979, Lee et al. 1980 et seq., Bond 1994, Logan et al. 1996, USFWS 2005); Pennsylvania (Smiley 1886, McDonald 1893, Hendricks et al. 1979, Cooper 1983, Pearson and Krumholz 1984, Tilmant 1999, Anonymous 2000, USFWS 2005); Rhode Island (Lapin, personal communication, Lee et al. 1980 et seq.); South Carolina (Baird 1887, McDonald 1886, 1893, Ravenel 1898, Hildebrand 1923, Dahlberg and Scott 1971a, 1971b, Loyacano 1975, Lee et al. 1980 et seq., Fretwell 2004); South Dakota (Anonymous 1892, Worth 1895, Cleary 1956, Shields 1958a, 1958b, Moyle and Clothier 1959, Underhill 1959, Bailey and Allum 1962, Lee et al. 1980 et seq., Young et al. 1997, USFWS 2005); Tennessee (McDonald 1886, Baird 1887, Anonymous 1892, Bean 1896, Ravenel 1896, Kuhne 1939, Ryon and Loar 1988, Etnier and Starnes 1993, Tilmant 1999, USFWS, Mississippi Museum of Natural Science 2004); Texas (Smiley 1886, Baird 1887, McDonald 1886, 1893, Bean 1896, Baughman 1950, Lee et al. 1980 et seq., Conner and Suttkus 1986, Howells 1992, Texas System of Natural Laboratories, Inc. and USGS 1994, Texas System of Natural Laboratories, Inc 1996, Red River Authority of Texas 2001, Texas Parks and Wildlife Department 1993, 1994, 2001, USFWS 2005, Anonymous 1994); Utah (Tanner 1936, Sigler and Miller 1963, Vanicek et al. 1970, Lee et al. 1980 et seq., Sigler and Sigler 1996, Tilmant 1999); Vermont (Countryman 1975, Lee et al. 1980 et seq.); Virginia (Smiley 1886, Baird 1887, 1889, McDonald 1886, 1893, Anonymous 1892, Worth 1895, Bean 1896, Ravenel 1896, 1898, Lee et al. 1980 et seq., Jenkins and Burkhead 1994, Tilmant 1999); Washington (Smith 1896, Chapman 1942, Lampman 1946, Wydoski and Whitney 1979, Lee et al. 1980 et seq., Beecher and Fernau 1982, Wydoski and Whitney 2003, USFWS 2005, Four Seasons Campground and Resort 2003); West Virginia (Clay 1962, Lee et al. 1980 et seq., Stauffer et al. 1995, USFWS 2005); Wisconsin (Johnson and Becker 1980, Becker 1983, Emery 1985, Burr and Page 1986, Fago 1992, Tilmant 1999, Jansen 2003, USFWS 2005); and Wyoming (Baxter and Simon 1970, Stone 1995)."

"Common carp has also been collected the Cidra, Guajataca, and Loiza reservoirs and the Lajas Irrigation Canal in Puerto Rico (Felix Grana, personal communication)."

Means of Introductions in the United States

From Nico et al. (2014):

“There is some question as to when and where common carp was first introduced into the United States. DeKay (1842) reported that the species was first brought into the United States from France by Henry Robinson of Orange County, New York in 1831 and 1832. In a letter to DeKay, Robinson detailed that he kept the fish in ponds and for several years released one to two dozen carp during the spring in the Hudson River near his residence, thereby creating a commercial fishery for the species. S. F. Baird of the U.S. Fish Commission examined fish taken from the Hudson River, as well as area fish then being sold on the New York markets, and reported that they were goldfish or goldfish hybrids and not true common carp (Redding 1884, Cole 1905). Whitworth (1996) cited early literature indicating common carp had been introduced into Connecticut as early as the 1840s; however, we question the positive identity of the species. Smith (1896) reported that common carp first appeared in the United States in 1872 when J. A. Poppe of Sonoma, California, imported five specimens from Germany and propagated them in private ponds for commercial purposes, mainly distributing them to applicants as a food fish (Smith 1896, Lampman 1946). In 1877, the U.S. Fish Commission imported common carp from Germany and for the next two decades the agency began stocking and distributing the species as food fish throughout much of the United States and its territories (Smiley 1886, Smith 1896, Cole 1905). State fish commissions also were commonly involved in distributing the species (e.g., Johnson and Becker 1980). Records from the early 1880s indicate that common carp stocked in farm ponds frequently escaped into open waters as a result of dam breaks or flood events (Smiley 1886). By 1885, the U.S. Fish Commission was actively stocking lakes and rivers throughout the country, often the fish were released from railroad tank cars at bridge crossing directly into streams (e.g., McDonald 1886). As a result of subsequent population growth and dispersal, common carp spread even further. More recently introductions of common carp have resulted because of the use of juvenile carp as bait fish (e.g., Swift et al. 1977). Various unusual genetic strains of common carp have been introduced into open waters the United States. In addition to the normal scaled carp, the U.S. Fish Commission distributed both mirror carp and leather carp varieties in the late 1800s (Smiley 1886, Cole 1905). Colorful varieties of common carp (i.e., nishikigoi or koi) are kept as pets in garden ponds and some have been introduced to ponds and public water bodies (Balon 1995). However, only a small percentage of common carp records in U.S. open waters are based on koi. Another cultured variety occasionally found in open waters is the Israeli carp (Robison and Buchanan 1988). Their presence in South Florida is believed to be the result of released bait with this species as a contaminant.”

Remarks

From Nico et al. (2014):

“Carp is only established in the Florida panhandle. It does not appear to be established in South Florida.”

“Balon (1995) reviewed the origin and history of domestication of common carp in Europe and elsewhere. Several agents of the U.S. Fish Commission documented the early years of common carp propagation and stocking in the United States (e.g., Smiley 1886, Smith 1896, Cole 1905). Although this species was popular in the early 1870s as a food fish, common carp fell into wide

disfavor soon after and is now considered a nuisance fish because of its abundance and detrimental effects on aquatic habitats. Trautman (1981) found common carp most abundant in streams enriched with sewage or with substantial runoff from agricultural land, but he reported it to be rare or absent in clear, cold waters, and streams of high gradient. Pflieger (1997) reported that the total weight and value of common carp taken by commercial fishermen in Missouri exceeded that of any other fish. Hartel et al. (1996) noted that more than 20,000 common carp were killed by a bacterial disease over a short period of time in the Merrimack River in the late 1970s. Because common carp have a higher salinity tolerance than most freshwater fishes, Swift et al. (1977) hypothesized that it may be spreading from one coastal stream to another through fresh or nearly fresh coastal waters in the Gulf area during periods of heavy rainfall and run-off, periods when salinities are greatly reduced.”

“DeVaney et al. (2009) performed ecological niche modeling to examine the invasion potential for common carp and three other invasive cyprinids (grass carp *Ctenopharyngodon idella*, black carp *Mylopharyngodon piceus*, and tench *Tinca tinca*). The majority of the areas where common carp have been collected, stocked, or have become established had a high predicted ecological suitability for this species.”

2 Biology and Ecology

Taxonomic Hierarchy and Taxonomic Standing

From ITIS (2011):

“Kingdom Animalia
 Subkingdom Bilateria
 Infrakingdom Deuterostomia
 Phylum Chordata
 Subphylum Vertebrata
 Infraphylum Gnathostomata
 Superclass Osteichthyes
 Class Actinopterygii
 Subclass Neopterygii
 Infraclass Teleostei
 Superorder Ostariophysi
 Order Cypriniformes
 Superfamily Cyprinoidea
 Family Cyprinidae
 Genus *Cyprinus*
 Species *Cyprinus carpio* Linnaeus, 1758

Taxonomic Status: Valid.”

Size, Weight, and Age Range

From Kottelat and Freyhof (2007):

“Maturity: Lm 35.1, range 25 - 36 cm; Max length : 110 cm SL male/unsexed; common length: 31.0 cm TL male/unsexed; (Chugunova 1959); max. published weight: 40.1 kg (Machacek 2007); max. reported age: 38 years (Hinton 1962).”

From Koch (2014):

A 64-year-old common carp was captured in Lake Gervais, Little Canada, Minnesota. The mean age of captured carp (n=127) was 31 years.

Environment

From Kottelat and Freyhof (2007):

“Freshwater; brackish; benthopelagic; pH range: 7.0 - 7.5; dH range: 10 - 15; potamodromous (Riede 2004).”

Climate/Range

From Kottelat and Freyhof (2007):

“Subtropical; 3°C - 35°C (Eaton et al. 1995); 60°N - 40°N.”

Distribution Outside the United States

Native

From Kottelat and Freyhof (2007):

“Europe to Asia: Black, Caspian and Aral Sea basins. Introduced throughout the world. Wild stocks are only present naturally in rivers draining to the Black, Caspian and Aral Sea. A reophilic wild population in the Danube is assumed to be the origin of the European species; this population is now under threat (Kottelat 1997).”

Introduced

From Kottelat and Freyhof (2007):

This species is reported as introduced in Thailand, Hawaii, Taiwan, Belarus, Indonesia, Albania, Belgium, Denmark, France, Germany, Greece, Ireland, Italy, Liechtenstein, Luxembourg, Malta, Netherlands, Norway, Portugal, Spain, Sweden, Switzerland, Bosnia Herzegovina, Botswana, Brunei, Quebec, Croatia, Czechoslovakia, Ecuador, French Polynesia, Guam, Haiti, Hong Kong, Jamaica, Kuwait, Kyrgyzstan, Lebanon, Lesotho, Latvia, Lithuania, Mozambique, Namibia, New Caledonia, Paraguay, Azores Is, East Timor, Reunion, Saudi Arabia, Singapore, Slovenia, Swaziland, Syria, Tajikistan, Togo, Turkey, Tanzania, Israel, Jordan River, Afghanistan, Myanmar, Iran, Nepal, Burundi, Argentina, Malaysia, Mexico, New Zealand, Poland, Romania, UK, Canada, Uruguay, South Africa, Mitidja Oueds, Finland, Australia, Chile, Estonia, Murgab

(Turkmenistan), British Columbia, Brazil, Japan, Kazakhstan (Lake Balkhash), Kenya, Sri Lanka, Madagascar, Philippines, Morocco, Zimbabwe, Cuba, Egypt, India, Colombia, Venezuela, Bolivia, Peru, Zambia, Congo, Dominican Republic, Puerto Rico, Guatemala, Nigeria, Iraq, Honduras, China, Papua New Guinea, Bangladesh, Rwanda, Ghana, Uganda, Nicaragua, Pakistan, El Salvador, Lao PDR, Tunisia, Central African Republic, Cyprus, Suriname, Cambodia, Viet Nam, Cameroon, Malawi, Korea, Sudan, Costa Rica, Hungary, Côte d'Ivoire, Mauritius, Panama, Former USSR, Angola (Pangula Lake), and Bhutan (FAO 1997).

From GISD (2014):

“*C. carpio* is the third most frequently introduced species world-wide (Welcomme 1992, in Saikia and Das 2009).”

Means of Introduction Outside the United States

From Kottelat and Freyhof (2007):

Reasons given for the introduction of this species include aquaculture, fisheries, diffused from other countries, angling, snail control, research, ornamental, accidental, weed control and unknown (FAO 1997). The species is reported to be established and having negative impacts on native populations in some areas where introduced (FAO 1997).

Short description

From Kottelat and Freyhof (2007):

“Dorsal spines (total): 3 - 4; Dorsal soft rays (total): 17-23; Anal spines: 2-3; Anal soft rays: 5 - 6; Vertebrae: 36 - 37. Diagnosed from other cyprinid species in Europe by having the following characters: 2 pairs of barbels; dorsal fin with 15-20½ branched rays; caudal fin deeply emarginate. Pharyngeal teeth 1, 1, 3:3, 1,1, robust, molar-like with crown flattened or somewhat furrowed. Scales large and thick. ‘Wild carp’ is generally distinguished by its less stocky build with height of body 1:3.2-4.8 in standard length. Very variable in form, proportions, squamation, development of fins, and color. Caudal fin with 3 spines and 17-19 rays (Spillman 1961). Last simple anal ray bony and serrated posteriorly; 4 barbels; 17-20 branched dorsal rays; body grey to bronze (Kottelat 2001). Also Suzuki and Yamaguchi 1980, Suzuki and Yamaguchi 1984.”

Biology

From Kottelat and Freyhof (2007):

“Adults inhabit warm, deep, slow-flowing and still waters such as lowland rivers and large, well vegetated lakes. Hardy and tolerant of a wide variety of conditions but generally favor large water bodies with slow flowing or standing water and soft bottom sediments. Thrive in large turbid rivers (Scott and Crossman 1973). Most active at dusk and dawn. Both adults and juveniles feed on a variety of benthic organisms and plant material. Spawns along shores or in backwaters. Adults often undertake considerable spawning migration to suitable backwaters and flooded meadows. Larvae survive only in very warm water among shallow submerged

vegetation. River regulation and hybridization with domesticated stocks, East Asian congeners and their hybrids have caused continuous decline of wild populations.”

Human uses

From Kottelat and Freyhof (2007):

“Fisheries: highly commercial; aquaculture: commercial; gamefish: yes; aquarium: commercial.”

“Utilized fresh and frozen (Frimodt 1995). Aquarium keeping: in groups of 5 or more individuals; minimum aquarium size >200 cm; not recommended for home aquariums (BMELF 1999).”

Diseases

From Kottelat and Freyhof (2007):

Water mold Disease (l.), Fungal diseases; Fin-rot Disease (late stage), Bacterial diseases; Red spot Disease, Bacterial diseases; Anchor worm Disease, Parasitic infestations; SVC, Viral diseases; Fish louse Infestation 1, Parasitic infestations; Coccidiosis (intestine), Parasitic infestations; Boil Disease, Parasitic infestations; Costia Disease, Parasitic infestations; Dactylogyrus Gill Flukes Disease, Parasitic infestations; Fish leech Infestation, Parasitic infestations; Trichodinosis, Parasitic infestations; Lymphocystis Disease, Viral diseases; Skin Flukes, Parasitic infestations; Fungal Gill Rot (sanguinis), Fungal diseases; Chilodonella Disease, Parasitic infestations; Worm Cataract, Parasitic infestations; Fin Rot (early stage), Bacterial diseases; Lymphocystis Disease (dark), Viral diseases; White spot Disease, Parasitic infestations; Trichodinosis, Parasitic infestations; Skin Flukes, Parasitic infestations; Trichodinella Infection 1, Parasitic infestations; Trichodina Infection 1, Parasitic infestations; Trichodina Infection 2, Parasitic infestations; Trichodina Infection 3, Parasitic infestations; Trichodinella Infection 2, Parasitic infestations; Myxobolus Infection 3, Parasitic infestations; Myxobolus Infection 4, Parasitic infestations; Trichodina Infection 5, Parasitic infestations; Myxobolus Infection 1, Parasitic infestations; Trichodina Infection 7, Parasitic infestations; Trichodinella Infection 3, Parasitic infestations; Thelohanellus Infection 1, Parasitic infestations; Anchorworm Disease (*Lernaea* sp.), Parasitic infestations; Dactylogyrus Gill Flukes Disease, Parasitic infestations; Trichodinosis, Parasitic infestations; Turbidity of the Skin (Freshwater fish), Parasitic infestations; Pallisentis Disease, Parasitic infestations; Columnaris Disease (l.), Bacterial diseases; Aeromonosis, Bacterial diseases; Infectious ascites (Ornament.), Bacterial diseases; Columnaris Disease (e.), Bacterial diseases; Unspecified tumors, Neoplasia (tumors of unknown origin); Hole-in-the-Head Disease, Parasitic infestations; Turbidity of the Skin (Freshwater fish), Parasitic infestations; Bacterial Infections (general), Bacterial diseases; Koi Herpes Virus, Viral diseases; Fish Pox Disease, Viral diseases; Congenital Deformities, Others; Carp Iridovirus, Viral diseases; Carp Coronavirus Infection, Viral diseases; Carp Reovirus, Viral diseases; Enteric Redmouth Disease, Bacterial diseases; Edwardsiellosis, Bacterial diseases; Epitheliocystis, Bacterial diseases; Capillaria Infestation 3, Parasitic infestations; Pseudocapillaria Infestation 1, Parasitic infestations; Myxobolus Infection 3, Parasitic infestations; Myxobolus Infection 3, Parasitic infestations; Anchor worm Disease, Parasitic infestations; Myxobolus Infection 3, Parasitic infestations and Velvet Disease 2 (*Piscinoodinium* sp.), Parasitic infestations.

SVC, Viral Diseases and Koi Herpes Virus are OIE-reportable.

Threat to humans

Potential pest.

3 Impacts of Introductions

From Nico et al. (2014):

“The common carp is regarded as a pest fish because of its widespread abundance and because of its tendency to destroy vegetation and increase water turbidity by dislodging plants and rooting around in the substrate, causing a deterioration of habitat for species requiring vegetation and clean water (Cole 1905, Cahoon 1953, Bellrichard 1996, Laird and Page 1996). Available literature indicates common carp may destroy aquatic macrophytes directly by uprooting or consuming the plants, or indirectly by increasing turbidity and thereby reducing light for photosynthesis. Bellrichard (1996) found that alterations in macrophyte biomass are due more to direct effects of common carp. In their review of the literature, Richardson et al. (1995) concluded that common carp has had noted adverse effects on biological systems including destruction of vegetated breeding habitats used by both fish and birds, and an increase in turbidity. It stirs up the bottom during feeding, resulting in increased siltation and turbidity (Lee et al. 1980 et seq.). This feeding behavior also destroys rooted aquatic plants that provide habitat for native fish species and food for waterfowl (Dentler 1993). There is also evidence that common carp prey on the eggs of other fish species (Moyle 1976, Taylor et al. 1984, Miller and Beckman 1996). For this reason, it may be responsible for the decline of the razorback sucker *Xyrauchen texanus* in the Colorado River basin (Taylor et al. 1984). In another case, Miller and Beckman (1996) documented white sturgeon *Acipenser transmontanus* eggs in the stomachs of common carp in the Columbia River. In California, carp have been implicated in the decrease in water clarity in Clear Lake, Lake County, and in the gradual disappearance of native fishes (Moyle 1976). McCarraher and Gregory (1970) wrote that in 1894 there was documentation that Sacramento perch *Archoplites interruptus* were becoming more scarce because carp was destroying their spawning grounds. Laird and Page (1996) stated that common carp may compete with ecologically similar species such as carpsuckers and buffalos. Because this species has been present in many areas since the first surveys, its impacts on many of the native fishes are difficult to determine. Once established in a water body, common carp is difficult and expensive to eliminate (e.g., Cahoon 1953).”

From GISD (2014):

“This species has been nominated as among 100 of the “World’s Worst” invaders.”

“*C. carpio* is the third most frequently introduced species world-wide (Welcomme 1992 in Saikia and Das 2009). On every continent where it has been introduced it has reduced water quality and degraded aquatic habitats (McCrimmon 1968, Roberts et al. 1995, King et al. 1997, [and] Koehn et al. 2000, in Jones and Stuart 2006).”

“Ecosystem Change: In shallow aquatic ecosystems, common carp can be considered “ecosystem engineers” or “keystone modifiers” (Jones et al. 1994, [and] Mills et al. 1993 in Parkos [et al.] 2003) in that they have strong effects on benthic communities. Aquatic macrophytes are integral to ecosystem functioning (Stansfield et al. 1997 in Nunn et al. 2007). Carp are known to damage aquatic macrophytes. [Macroinvertebrates] are keystone species in aquatic ecosystems (Scheffer 1998, [and] Scheffer et al. 2001 in Shin-ichiro et al. 2009). Shin-ichiro and colleagues (2009) found carp significantly influenced benthic macroinvertebrates.”

This type of impact has been reported from Argentina (Ref. 1739 in FishBase 2009), Australia (NSW Department of Primary Industries 2005), Canada (FishBase 2009), Chile (FishBase 2009), Republic of the Congo (FishBase 2009), Cyprus (FishBase 2009), India (FishBase 2009), Kenya (FishBase 2009, Britton et al. 2007), Madagascar (FishBase 2009), Malawi (FishBase 2009), Morocco (FishBase 2009), New Zealand (FishBase 2009, DOC Undated), Papua New Guinea (Coates and Ulaivi 1995, FishBase 2009), Singapore (FishBase 2009), Switzerland (FishBase 2009), Tunisia (FishBase 2009), and Uruguay (FishBase 2009).

“Habitat Alteration: Carp may pose a threat to wetlands that are used by many fish as spawning and nursery habitats (Parkos [et al.] 2003).”

This type of impact has been reported from Australia (Roberts et al. 1995, Jones and Stuart 2009), China (Jia et al. 2008), Mexico (Zambrano et al. 1999, Hincosa-Garro and Zambrano 2004), Poland, Portugal (FishBase 2009), the United Kingdom (Gordon Copp personal communication 2006 in FishBase 2009), and the United States (Hutchinson 1975, Lodge 1991, Newman 1991, Cowl et al. 1998, and Persson and Crowder 1998 in Miller and Provenza 2007; Miller and Provenza 2007).

“Modification of natural benthic communities: Carp are believed to stimulate algal bloom formation by increasing nutrient release from sediments and decreasing algal grazing by cladocerans (which the juvenile carp prey upon) (Pinto et al. 2005).”

This type of impact has been reported from Australia (Harris 1995 in Pinto et al. 2005) and the United States.

“Modification of nutrient regime: Carp increase nutrients in the water column in two ways: by sediment resuspension and by excretion (Lamarra 1975 [and] Brabrand et al. 1990 in Chumchal 2002).”

This type of impact has been reported from Australia (NSW Department of Primary Industries 2005, Pinto et al. 2005, and King et al. 1997 in Pinto et al. 2005), Canada (Lougheed et al. 1998 in Pinto et al. 2005), and the United States (Lougheed et al. 1998 in Pinto et al. 2005).

“Reduction in native biodiversity: In California, USA, carp have been implicated in the gradual disappearance of native fishes (Moyle 1976a in Nico [et al.] 2009). Data from Miller and Crowl (2006) suggests that carp can significantly affect species abundance and diversity of macrophytes and some macroinvertebrates. Common carp negatively affected macrophyte abundance by reduction of light availability, increase of siltation rates, ingestion of plant matter and uprooting

during feeding activity (Parkos [et al.] 2003). The loss of rooted macrophytes due to carp activity is intuitively likely to lead to a decline in biological diversity, in endemic fish, amphibians, and reptiles in Mexico (Crowder and Painter 1991 in Zambrano et al. 1999) and elsewhere.”

This type of impact has been reported from Argentina (Ref. 1739 in FishBase 2009), Australia (Fletcher et al. 1985, King and Hunt 1967, Driver et al. 1997, and Roberts 1995 in Pinto et al. 2005; Fletcher et al. 1985; NSW Department of Primary Industries 2005; South Australia Government Undated), Canada (Refs 1998 and 1995 in FishBase 2009, Loughheed et al. 1998 in Pinto et al. 2005), Kenya (Hickey and Harper 2002 in Hickley et al. 2004, Hickley et al. 2004, FishBase 2009), Mexico (Ref 6447 in FishBase 2009, Hinojosa-Garro and Zambrano 2004), New Zealand (Ref 13730 in FishBase 2009), Papua New Guinea (Coates and Ulaiwi 1995), the United States (Loughheed et al. 1998 in Pinto et al. 2005, Heckman et al. 1981, National Archives Microfilm Publications 1850-1902, and Brotherson 1981 in Miller and Provenza 2007; Miller and Provenza 2007), and Venezuela (FishBase 2009).

“Physical disturbance: Carp stir up bottom sediments during feeding, resulting in increased siltation and bioturbation (Arlinghaus and Mehner 2003; Parkos [et al.] 2003; Lee et al. 1980 in Nico Maynard and Schofield 2009).”

This type of impact has been reported from Australia (NSW Department of Primary Industries 2005), India (FishBase 2009), Kenya (Petr 2000 in Hickley et al. 2004), and the Netherlands (Meijer et al. 1990 in Hickley et al. 2004).

“Threat to endangered species: Non-native fish can drive native species to local extinction (Zambrano et al. 2006). Predation: Carp prey on macroinvertebrates (Parkos [et al.] 2003). There is also evidence that common carp prey on the eggs of other fish species (Moyle 1976a, Taylor et al. 1984, [and] Miller and Beckman 1996 in Nico Maynard and Schofield 2009).”

“Competition: Laird and Page (1996, in Nico [et al.] 2009) stated that common carp may compete with ecologically similar species such as carp suckers and buffalo fish.”

This type of impact has been reported from Australia (Adamek 1998, Koehn et al. 2000, and Koehn & O'Connor 1990a in Koehn 2004; FishBase 2009; McDowall 1996 and Mallen-Cooper and Stuart 2003 in Jones and Stuart 2009; Nicol et al. 2004 and Koehn et al. 2000a in Nicol et al. 2004; Nicol et al. 2004; Jones and Stuart 2006), China (Jia et al. 2008), India (FishBase 2009), Madagascar (FishBase 2009), Mexico (Hinojosa-Garro and Zambrano 2004), Papua New Guinea (Ref 2847 in FishBase 2009), and Poland.

“Economic/Livelihoods: Growth rates and stocks of other fish may be impacted by competition with carp (Arlinghaus and Mehner 2003), including perch. Carp provide an important source of protein in some third world countries (FishBase 2009).”

This type of impact has been reported from Algeria (FishBase 2009), Argentina (FishBase 2009), Australia, Belgium (FishBase 2009), Bolivia, Cambodia (FishBase 2009), the Republic of the Congo (FishBase 2009), Costa Rica (FishBase 2009), Croatia (FAO 1997 in FishBase 2009), Cyprus (FishBase 2009), Estonia (FishBase 2009), Hungary (FishBase 2009), India (FishBase

2009), Indonesia (FishBase 2009), the Islamic Republic of Iran (FishBase 2009), Iraq (FishBase 2009), Kenya (FishBase 2009), Kyrgyzstan (FishBase 2009), the Lao People's Democratic Republic (FishBase 2009), Lesotho (FishBase 2009), Madagascar (FishBase 2009), Malawi (FishBase 2009), Malaysia (FishBase 2009), Mexico (FishBase 2009, Zambrano et al. 1999), Morocco (FishBase 2009), Mozambique (FishBase 2009), Myanmar (FishBase 2009), Namibia (FishBase 2009), Nepal (FishBase 2009), Netherlands (FishBase 2009), Nicaragua (FishBase 2009), Pakistan (FishBase 2009), Papua New Guinea (Coates and Ulaiwi 1995, Coates 1993b in Coates and Ulaiwi 1995), Paraguay (FishBase 2009), Poland (FishBase 2009), Puerto Rico (FishBase 2009), Romania (FishBase 2009), South Africa (FishBase 2009), Switzerland (FishBase 2009), Thailand, Turkmenistan (FishBase 2009), the United Kingdom (Nunn et al. 2007), and Vietnam (FishBase 2009).

“Human nuisance: By stirring up river substrate and reducing aquatic vegetation carp can makes waterways unattractive and can render the water unsuitable for swimming or for drinking by livestock (NIWA 2003).”

Disease transmission

This type of impact has been reported from Australia (Environment ACT Undated, NSW Department of Primary Industries 2005), Mexico (Ref 48731 in FishBase 2009), and South Africa.

Hybridization

This type of impact has been reported from Turkmenistan (Ref 30700 in FishBase 2009).

For additional information on location-specific impacts, please see GISD (2014).

From Váradi (2014):

“*C. carpio* has been used in aquaculture almost throughout human history, being cultured in China since at least 475 BC. Its ornamental forms in Japan (koi) are still symbolic of courage and energy. *C. carpio* is a native of Asia, from where it was spread by humans through Europe, and is now established on all continents except Antarctica – it can be considered the world's most widely distributed freshwater fish. It is important in many parts of the world and continues to be used both in pond and captive fisheries because of its potentially rapid growth in eutrophic waters and ability to tolerate adverse environmental conditions. *C. carpio* is also highly prized by anglers in many countries. However it is considered as a pest in North America, Australia and New Zealand (Dowal 1996). As a zooplankton feeder in the juvenile stage and a benthic feeder later on, *C. carpio* may contribute considerably to algal (including cyanobacterial) blooms.”

“Adverse impacts of carp on the natural water ecosystems have been proved in some places. Carp have uprooted aquatic plants and destroyed the aquatic weed populations (Laird and Page 1996). They also increase the turbidity of waters by digging and stirring the upper layer of the bottom. As a consequence of this, light penetration decreases, which destroys the macrophyte populations in the spawning areas of phytophil species. Carp may directly consume the eggs of

other species (Miller and Beckman 1966). By destroying the spawning substrata, by consumption of eggs of native species (Tyus and Saunders 2000) and by competing with species having similar feeding habit (Laird and Page 1995) the presence of carp may be one reason why the populations of native species are declining.”

Parkos et al. (2003):

“We examined the effects of adult common carp (*Cyprinus carpio*) on shallow aquatic ecosystems and compared the effects with those of a native benthic fish, channel catfish (*Ictalurus punctatus*). Experimental ponds contained enclosures (0.06 ha) with a low carp biomass (174 kg·ha⁻¹), high carp biomass (476 kg·ha⁻¹), high catfish biomass (416 kg·ha⁻¹), and no fish. We measured abiotic factors (turbidity, suspended solids, total phosphorus), as well as effects on adjacent trophic levels (aquatic macrophytes, zooplankton, and aquatic macroinvertebrates) from July to September. Common carp was positively related to total phosphorus, turbidity, suspended solids, and zooplankton biomass, and negatively related to macrophyte and macroinvertebrate abundance. Suspended solids in the carp treatments consisted primarily of inorganic particles. Carp were either positively or negatively related to phytoplankton, depending on zooplankton abundance. A high biomass of carp had greater effect on nutrients, turbidity, and suspended solids than a low biomass. Channel catfish was positively related to total phosphorus concentrations and altered zooplankton composition, but did not affect turbidity, suspended solids, macroinvertebrates, and macrophytes. These results suggest that common carp have a stronger influence on water quality and aquatic community structure than benthic fish native to North America.”

From Weber and Brown (2009):

“Common carp in shallow lakes increase turbidity, phytoplankton, and blue-green algae; mobilize nutrients; and decrease macrophytes, macroinvertebrates, and fishes (Lamarra 1975, Parkos et al. 2003, Egertson and Downing 2004). Common carp populations function as ecosystem engineers by regulating bottom-up and top-down processes. Introduced common carp are not regulated by topdown processes (i.e., experience minimal predation) and are not limited by bottom-up processes (i.e., prey resources); thus, common carp can regulate ecosystems through a middle-out framework. Ecosystem effects imposed by common carp may be greater than those caused by other native benthivores and occur even at relatively low common carp densities, with increases in undesirable ecosystem effects with increases in common carp biomass (Parkos et al. 2003, Chumchal et al. 2005). Ecosystem engineering by common carp can transform lakes from clear to turbid water, with ecosystem-wide consequences (Scheffer 1998, Haas et al. 2007).”

4 Global Distribution




Figure 1. Global distribution of *Cyprinus carpio*. Map from GBIF (2011).

5 Distribution within the United States




Figure 2. U.S. distribution of *Cyprinus carpio*. Map from Nico et al. (2014).

6 CLIMATCH

Summary of Climate Matching Analysis

The climate match (Australian Bureau of Rural Sciences 2008; 16 climate variables; Euclidean Distance) was high across the entire contiguous U.S. Climate 6 proportion indicated that the contiguous U.S. has a high climate match. The range for a high climate match is 0.103 and greater; climate match of *Cyprinus carpio* is 0.999.




Figure 3. CLIMATCH (Australian Bureau of Rural Sciences 2008) source map showing weather stations selected as source locations (red) and non-source locations (blue) for *Cyprinus carpio* climate matching. Source locations from GBIF (2011) and Nico et al. (2014).




Figure 4. Map of CLIMATCH (Australian Bureau of Rural Sciences 2008) climate matches for *Cyprinus carpio* in the contiguous United States based on source locations reported by GBIF (2011) and Nico et al. (2014). 0= Lowest match, 10=Highest match.

Table 1. CLIMATCH (Australian Bureau of Rural Sciences 2008) climate match scores.

CLIMATCH Score	0	1	2	3	4	5	6	7	8	9	10
Count	0	0	1	0	0	1	11	45	197	113	1606
Climate 6 Proportion =		0.999 (High)									

7 Certainty of Assessment

The biology, distribution, and negative impacts of *Cyprinus carpio* are adequately documented in the scientific literature. No further information is needed to evaluate the negative impacts the species is having where introduced. Certainty of this assessment is high.

8 Risk Assessment

Summary of Risk to the Contiguous United States

Cyprinus carpio is a freshwater and brackish water fish native to the Caspian, Black, and Aral Sea basins. This species is now established in locations worldwide, including the U.S. Only Alaska has not had a documented introduction of this species, and the species has established in all other states except Maine. *Cyprinus carpio* is used for aquaculture, commercial fisheries, aquaria, and bait, which has promoted its spread. Negative impacts of this species are well-studied. The primary impacts of this species are habitat modification and reduction of the abundance of macrophytes and macroinvertebrates. When foraging for benthic organisms, this species uproots macrophytes and disturbs sediments. The resultant increase in turbidity and phosphorus loading promotes algal blooms, degrades water quality, prevents growth of macrophytes, and possibly causes effects on higher organisms. This species has also been implicated in the decline of native fish due to competition and egg predation. *Cyprinus carpio* is also a known carrier of OIE-reportable diseases. Climate match with the contiguous U.S. is high. Overall risk for this species is high.

Assessment Elements

- **History of Invasiveness (Sec. 3):** High
- **Climate Match (Sec.6):** High
- **Certainty of Assessment (Sec. 7):** High
- **Remarks/Important additional information** Host of 63 diseases/parasites some of which are OIE-reportable and listed as a potential pest.
- **Overall Risk Assessment Category: High**

9 References

Note: The following references were accessed for this ERSS. References cited within quoted text but not accessed are included below in Section 10.

- Australian Bureau of Rural Sciences. 2008. CLIMATCH. Available:
<http://data.daff.gov.au:8080/Climatch/climatch.jsp>. (April 2011).
- Frankis, M., V. Jacinto, and L. Shapiro. 2014. *Cyprinus carpio*. Encyclopedia of Life. Available:
<http://eol.org/pages/985921/overview>. Photo license available:
<http://creativecommons.org/licenses/by-nc/3.0/legalcode>. (September 2014).
- Froese, R., and D. Pauly. Editors. 2011. FishBase. Available:
<http://www.fishbase.us/summary/Cyprinus-carpio.html>. (April 2011).
- Global Biodiversity Information Facility (GBIF). 2011. Available:
<http://www.gbif.org/species/4286975>. (April 2011).
- Global Invasive Species Database (GISD). 2014. *Cyprinus carpio*. Available from:
<http://www.issg.org/database/species/ecology.asp?si=60&fr=1&sts=&lang=EN>. (July 2014).
- ITIS. 2011. *Cyprinus carpio*. Integrated Taxonomic Information System. Available:
http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=163344. (April 2011).
- Koch, J.D. 2014. Source-sink population structure of invasive common carp in a model Midwestern watershed: empirical evidence and notes on management. M. Sc. thesis. Department of Fisheries, Wildlife, and Conservation Biology, University of Minnesota, Saint Paul, Minnesota.
- Kottelat, M., and J. Freyhof. 2007. Handbook of European freshwater fishes. Publications Kottelat, Cornol, Switzerland.
- Nico, L., E. Maynard, P.J. Schofield, M. Cannister, J. Larson, A. Fusaro, and M. Neilson. 2014. *Cyprinus carpio*. USGS Nonindigenous Aquatic Species Database, Gainesville, Florida. Available: <http://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=4>. Revision Date: 6/26/2014.
- Parkos, J.J., III, V.J. Santucci, Jr., and D.H. Wahl. 2003. Effects of adult common carp (*Cyprinus carpio*) on multiple trophic levels in shallow mesocosms. Canadian Journal of Fisheries and Aquatic Sciences 60(2): 182-192.
- Váradi, L. 2014. *Cyprinus carpio* (Common Carp). CAB International, Wallingford, United Kingdom. Available: <http://www.cabi.org/isc/datasheet/17522>. (July 2014).

Weber, M.J., and M.L. Brown. 2009. Effects of common carp on aquatic ecosystems 80 years after “Carp as a dominant”: Ecological insights for fisheries management. *Reviews in Fisheries Science* 17(4): 524-537.

10 References Quoted But Not Accessed

Note: The following references are cited within quoted text within this ERSS, but were not accessed for its preparation. They are included here to provide the reader with more information.

American Fisheries Society, Idaho Chapter. 2001. Fish ID. Available:
<http://www.fisheries.org/idaho/>.

Anonymous. 1892. Report of distribution of fish and eggs from July 1, 1888, to June 30, 1889. Pages 379-394 *in* Report of the Commissioner for 1888. Part XVI. U.S. Commission of Fish and Fisheries, Washington, District of Columbia.

Anonymous. 1994. Water body records. *Texas Parks & Wildlife News* 1994(February 25): 1-19.

Anonymous. 2000. Northwestern PA waters. *James's Northeastern Fishing Guide*.

Anonymous. 2001. Fishing Records - Nevada. *Insider Viewpoint Magazine*.

Arlinghaus, R., and T Mehner. 2003. Socio-economic characterization of specialized common carp (*Cyprinus carpio* L.) anglers in Germany, and implications for inland fisheries management and eutrophication control. *Fisheries Research* 61: 19-33.

Bailey, R.M., and M.O. Allum. 1962. Fishes of South Dakota. Miscellaneous Publication No. 119 of the Museum of Zoology, University of Michigan, Ann Arbor, Michigan.

Bailey, R.M., and G.R. Smith. 1981. Origin and geography of the fish fauna of the Laurentian Great Lakes basin. *Canadian Journal of Fisheries and Aquatic Sciences* 38: 1539-1561.

Baird, S.F. 1887. Report of the Commissioner for 1885. Part XIII. U.S. Commission of Fish and Fisheries, Washington, District of Columbia.

Baird, S.F. 1889. Report of the Commissioner for 1886. Part XIV. U.S. Commission of Fish and Fisheries, Washington, District of Columbia.

Balon, E.K. 1995. Origin and domestication of the wild carp, *Cyprinus carpio*: from Roman gourmets to the swimming flowers. *Aquaculture* 129: 3-48.

Baughman, J. L. 1950. Random notes on Texas fishes. *Texas Journal of Science* 2: 117-138.

Baxter, G.T., and J.R. Simon. 1970. Wyoming fishes. Wyoming Game and Fish Department Bulletin 4, Cheyenne, Wyoming.

- Bean, T.H. 1903. Catalogue of the Fishes of New York. New York State Museum Bulletin 60, Zoology 9. University of the State of New York Bulletin 278.
- Becker, G.C. 1983. Fishes of Wisconsin. University of Wisconsin Press, Madison, Wisconsin.
- Beecher, H.A., and R.F. Fernau. 1982. Fishes of Oxbow Lakes of Washington. Northwest Science 57(2): 125-131.
- Bellrichard, S.J. 1996. Effects of common carp (*Cyprinus carpio*) on submerged macrophytes and water quality in a backwater lake on the upper Mississippi River. Master's thesis, University of Wisconsin-La Crosse. Reprinted by the National Biological Service, Environmental Management Technical Center, Onalaska, Wisconsin.
- Blatchley, W.S. 1938. The Fishes of Indiana: with Descriptions, Notes on Habits and Distribution in the State. The Nature Publishing Co., Indianapolis, Indiana.
- Bond, C.E. 1994. Keys to Oregon freshwater fishes. Oregon State University Bookstores, Inc., Corvallis. 58:1-42.
- Boschung, H.T. 1992. Catalogue of freshwater and marine fishes of Alabama. Alabama Museum of Natural History Bulletin 14: 1-266.
- Bozeman, L., and P. Charp. 2001. Army Materials Technology Lab, Watertown, Middlesex, Massachusetts.
- Bradley, W.G., and J.E. Deacon. 1967. The biotic communities of southern Nevada. Nevada State Museum Anthropological Papers 13(4): 201-273.
- Breukelman, J. 1946. A review of Kansas ichthyology. Transactions of the Kansas Academy of Science 49: 51-70.
- Britton, J.R., R.R. Boar, J. Grey, J. Foster, J. Lugonzo, and D.M. Harper. 2007. From introduction to fishery dominance: the initial impacts of the invasive carp *Cyprinus carpio* in Lake Naivasha, Kenya, 1999 to 2006. Journal of Fish Biology (2007) 71 (Supplement D): 239–257.
- Brock, V.E. 1960. The introduction of aquatic animals into Hawaiian waters. Internationale Revue der gesamten Hydrobiologie 454: 463-480.
- Brown, C.J.D. 1971. Fishes of Montana. Montana State University, Bozeman, Montana.
- Bundesministerium für Ernährung, Landwirtschaft und Forsten (BMELF). 1999. Gutachten über Mindestanforderungen an die Haltung von Zierfischen (Süßwasser). Bundesministerium für Ernährung, Landwirtschaft und Forsten (BMELF), Bonn, Germany.

- Burkhead, N.M., S.J. Walsh, B.J. Freeman, and J.D. Williams. 1997. Status and restoration of the Etowah River, an imperiled southern Appalachian ecosystem. Pages 375-444 in G.W. Benz, and D.E. Collins, editors. Aquatic Fauna in Peril: The Southeastern Perspective. Special Publication 1, Southeast Aquatic Research Institute, Lenz Design & Communications, Decatur, Georgia.
- Burr, B.M., and L.M. Page. 1986. Zoogeography of fishes of the lower Ohio-upper Mississippi basin. Pages 287-324 in C.H. Hocutt, and E.O. Wiley, editors. The Zoogeography of North American Freshwater Fishes. John Wiley and Sons, New York, New York.
- Burr, B.M., and M.L. Warren, Jr. 1986. A Distributional Atlas of Kentucky Fishes. Kentucky Nature Preserves Commission Scientific and Technical Series 4.
- Cahoon, W.G. 1953. Commercial carp removal at Lake Mattamuskeet, North Carolina. Journal of Wildlife Management 17(3): 312-317.
- Call, L.E. 1961. Agricultural research at Kansas State Agricultural College (KSU) before enactment of the Health Act (1887). Agricultural Experimental Station, Kansas State University, Bulletin 441: 1-43.
- Chapman, W.M. 1942. Alien fishes in the waters of the Pacific northwest. California Fish and Game 28: 9-15.
- Chugunova, N.I. 1959. Age and growth studies in fish. A systematic guide for ichthyologists. Israel Program for Scientific Translations. Izdatel'stvo Akademii Nauk SSSR, Moskva.
- Chumchal, M. 2002. *Cyprinus carpio*, Animal Diversity Web. Available from: http://animaldiversity.ummz.umich.edu/site/accounts/information/Cyprinus_carpio.html. (September 2009).
- Chumchal, M.M., W.H. Nowlin, and R.W. Drenner. 2005. Biomass dependent effects of carp on water quality in shallow ponds. Hydrobiologia 545: 271-277.
- Clay, W.M. 1962. Aquatic-life resources of the Ohio River. Ohio River Valley Water Sanitation Commission, Cincinnati, Ohio.
- Clay, W.M. 1975. The Fishes of Kentucky. Kentucky Department of Fish and Wildlife Resources, Frankfort, Kentucky.
- Cleary, R.E. 1956. The distribution of the fishes of Iowa. Pages 267-324 in J.R. Harlan, and E.B. Speaker, editors. Iowa fish and fishing. Iowa Conservation Commission, Des Moines, Iowa.

- Coates, D., and W.K. Ulaiwi. 1995. A simple model for predicting ecological impacts of introduced aquatic organisms: a case study of common carp, *Cyprinus carpio* L., in the Sepik-Ramu River Basin, Papua New Guinea. *Fisheries Management and Ecology* 2(3): 227-242.
- Cobb, J.N. 1902. Commercial fisheries of the Hawaiian Islands. Pages 381-499 *in* Report of the Commissioner for the year ending June 30, 1901. U.S. Commission of Fish and Fisheries, Government Printing Office, Washington, District of Columbia.
- Cole, L.J. 1905. The German carp in the United States. Pages 523-641 *in* Report of the Bureau of Fisheries for 1904. U.S. Department of Commerce and Labor. Government Printing Office, Washington, District of Columbia.
- Conner, J.V., and R.D. Suttkus. 1986. Zoogeography of freshwater fishes of the western Gulf Slope of North America. Pages 413-456 *in* C.H. Hocutt, and E.O. Wiley, editors. *The Zoogeography of North American Freshwater Fishes*. John Wiley and Sons, New York, New York.
- Cook, F.A. 1959. *Freshwater Fishes of Mississippi*. Mississippi Game and Fish Commission Jackson, Mississippi.
- Cooper, E.L. 1983. *Fishes of Pennsylvania and the Northeastern United States*. Pennsylvania State University Press University Park, Pennsylvania.
- Countryman, W.D. 1975. Checklist of the recent fishes of Vermont. Norwich University, Northfield, Vermont. Unpublished mimeograph (revised version, October 1975).
- Courtenay, W.R., Jr. 1970. Florida's walking catfish. *Ward's Natural Science Bulletin* 10(69): 1, 4, 6.
- Courtenay, W.R., Jr. 1985. Florida Atlantic University Quarterly Reports for 1985 to the U.S. Fish and Wildlife Service, Gainesville, Florida.
- Courtenay, W.R., Jr., H.F. Sahlman, W.W. Miley, II, and D.J. Herrema. 1974. Exotic fishes in fresh and brackish waters of Florida. *Biological Conservation* 6(4): 292-302.
- Cross, F.B. 1967. *Handbook of Fishes of Kansas*. State Biological Survey and University of Kansas Museum of Natural History, Miscellaneous Publication 45, Topeka, Kansas.
- Cross, F.B., and J.T. Collins. 1995. *Fishes in Kansas*. University of Kansas Natural History Museum.
- Cudmore-Vokey, B., and E.J. Crossman. 2000. Checklists of the fish fauna of the Laurentian Great Lakes and their connecting channels. *Canadian Manuscript Report of Fisheries and Aquatic Sciences* 2500.

- Dahlberg, M.D., and D.C. Scott. 1971a. The freshwater fishes of Georgia. Bulletin of the Georgia Academy of Science 29: 1-64.
- Dahlberg, M.D., and D.C. Scott. 1971b. Introductions of freshwater fishes in Georgia. Bulletin of the Georgia Academy of Science 29: 245-252.
- Deacon, J.E., and J.E. Williams. 1984. Annotated list of the fishes of Nevada. Proceedings of the Biological Society of Washington 97(1): 103-118.
- DeKay, J.E. 1842. Zoology of New-York, or the New-York fauna. Part IV. Fishes. W. and A. White and J. Visscher, Albany, New York.
- Dentler, J.L. 1993. Noah's farce: The regulation and control of exotic fish and wildlife. University of Puget Sound Law Review 17: 191-242.
- DeVaney, S.C., K.M. McNyset, J.B. Williams, A.T. Peterson, and E.O. Wiley. 2009. A tale of four "carp": invasion potential and ecological niche modeling. PLoS ONE 4(5): e5451.
- Devick, W.S. 1991. Patterns of introductions of aquatic organisms to Hawaiian freshwater habitats. Pages 189-213 *in* New directions in research, management and conservation of Hawaiian freshwater stream ecosystems. Proceedings of the 1990 symposium on freshwater stream biology and fisheries management, Division of Aquatic Resources, Hawaii Department of Land and Natural Resources.
- Department of Conservation (DOC), New Zealand. Undated. Conservation > Threats & impacts > Animal pests > Animal pests A - Z > Fish > Facts > Koi carp.
- Douglas, N.H., and J.T. Davis. 1967. A checklist of the freshwater fishes of Louisiana. Louisiana Wildlife and Fisheries Commission.
- Douglas, N.H. 1974. Freshwater Fishes of Louisiana. Claitor's Publishing Division, Baton Rouge, Louisiana.
- Dowal, R. 1996. Freshwater Fish of South-Eastern Australia. Reed Books, Chatswood, NSW, Australia.
- Dyche, L.L. 1914. Ponds, pond fish and pond fish culture. Kansas Department of Fish and Game, Bulletin 1: 1-208.
- Eaton, J.G., J.H. McCormick, B.E. Goodno, D.G. O'Brien, H.G. Stefany, M. Hondzo, and R.M. Scheller. 1995. A field information-based system for estimating fish temperature tolerances. Fisheries 20(4): 10-18.
- Eddy, S., and J.C. Underhill. 1974. Northern fishes, with special reference to the upper Mississippi Valley. 3rd Edition. University of Minnesota Press, Minneapolis, Minnesota.

- Egertson, C.J., and J.A. Downing. 2004. Relationship of fish catch and composition to water quality in a suite of agriculturally eutrophic lakes. *Canadian Journal of Fisheries and Aquatic Sciences* 61: 1784–1796.
- Ellis, M.M. 1974. *Fishes of Colorado*. University of Colorado Studies, Boulder, Colorado.
- Emery, L. 1985. Review of fish introduced into the Great Lakes, 1819-1974. Great Lakes Fishery Commission Technical Report 45: 1-31.
- Etnier, D.A., and W.C. Starnes. 1993. *The Fishes of Tennessee*. University of Tennessee Press, Knoxville, Tennessee.
- Evermann, B.W., and C. Rutter. 1895. The fishes of the Colorado Basin. *Bulletin of the U.S. Fish Commission for 1894*, 14: 473-486.
- Everhart, W.H. 1976. *Fishes of Maine*. Maine Department of Inland Fisheries and Wildlife. Augusta, Maine.
- Fago, D. 1992. Distribution and relative abundance of fishes in Wisconsin. VIII. Summary Report. Technical Bulletin, Department of Natural Resources, Madison, Wisconsin.
- FAO. 1997. *FAO Database on Introduced Aquatic Species*. FAO Database on Introduced Aquatic Species, FAO, Rome.
- Ferguson, T.B. 1876. Report of the Commissioners of Fisheries of Maryland to the General Assembly. 1 January 1876. John F. Wiley, Annapolis, Maryland.
- FishBase. 2009. *Cyprinus carpio carpio*. Common Names Summary: Available from: <http://www.fishbase.org/comnames/CommonNamesList.php?ID=1450&GenusName=Cyprinus&SpeciesName=carpio+carpio&StockCode=1643>. (September 2009).
- Fletcher, A.R., A.K. Morison, and D.J. Hume. 1985. Effects of Carp, *Cyprinus carpio* L., on Communities of Aquatic Vegetation and Turbidity of Waterbodies in the Lower Goulburn River Basin. *Australian Journal of Marine and Freshwater Research* 36: 311-27.
- Four Seasons Campground and Resort. 2003. Fishing Information. Available: <http://www.fourseasonscampground.com/fishinginfo.asp>.
- Fowler, H.W. 1906. The fishes of New Jersey. Pages 35-477 in *Annual Report of the New Jersey State Museum (1905)*, part II. MacCrellich and Quigley, State Province, Trenton, New Jersey.
- Fowler, H.W. 1952. A list of the fishes of New Jersey, with off-shore species. *Proceedings of the Academy of Natural Sciences of Philadelphia CIV*: 89-151.

- Fretwell, S. 2004. Thousands of carp dying in lakes. The State.com. May 26, 2004.
- Frimodt, C. 1995. Multilingual illustrated guide to the world's commercial warmwater fish. Fishing News Books, Osney Mead, Oxford, England.
- Gerking, S.D. 1945. Distribution of the fishes of Indiana. Investigations of Indiana Lakes and Streams 3: 1-137.
- Gilbert, C.H., and N.B. Scofield. 1898. Notes of a collection of fishes from the Colorado Basin in Arizona. Proceedings of the U.S. National Museum 20: 487-499.
- Haas, K., U. Kohler, S. Diehl, P. Köhler, S. Dietrich, S. Holler, A. Jaensch, M. Niedermaier, and J. Vilsmeier. 2007. Influence of fish on habitat choice of water birds: A whole system experiment. Ecology 88: 2915–2925.
- Hall, G.E. 1956. Additions to the fish fauna of Oklahoma with a summary of introduced species. Southwestern Naturalist 1(1): 16-26.
- Harlan, J.R., E.B. Speaker, and J. Mayhew. 1987. Iowa fish and fishing. Iowa Department of Natural Resources, Des Moines, Iowa.
- Hartel, K.E. 1992. Non-native fishes known from Massachusetts freshwaters. Occasional Reports of the Museum of Comparative Zoology, Harvard University, Fish Department, Cambridge, Massachusetts.
- Hartel, K.E., D.B. Halliwell, and A.E. Launer. 1996. An annotated working list of the inland fishes of Massachusetts, University of Massachusetts, Cambridge, Massachusetts.
- Hay, O.P. 1894. The lampreys and fishes of Indiana. Report of the State Geologist.
- Hendricks, M.L., J.R. Stauffer, Jr., C.H. Hocutt, and C.R. Gilbert. 1979. A preliminary checklist of the fishes of the Youghiogheny River. Chicago Academy of Sciences, Natural History Miscellanea 203: 1-15.
- Hickley, P., S.M. Muchiri, J.R. Britton, and R.R. Boar. 2004. Discovery of carp, *Cyprinus carpio*, in the already stressed fishery of Lake Naivasha, Kenya. Fisheries Management & Ecology 11(2): 139-142.
- Hildebrand, S.F. 1923. Annotated list of fishes collected in the vicinity of Augusta, Georgia, with a description of a new darter. Bulletin of the U.S. Bureau of Fisheries 39: 1-8.
- Hinojosa-Garro, D., and L. Zambrano. 2004. Interactions of common carp (*Cyprinus carpio*) with benthic decapods in shallow ponds. Hydrobiologia 515(1-3): 115-122.
- Hinton. 1962. Horned shark, gar, mormyriad, characin, carp, armored catfish, arowana, upside down catfish. Not given.

- Holton, G.D. 1990. A Field Guide to Montana Fishes. Montana Department of Fish, Wildlife and Parks Helena, Montana.
- Howells, R.G. 1992. Annotated list of introduced non-native fishes, mollusks, crustaceans and aquatic plants in Texas waters. Texas Parks and Wildlife Department, Management Data Series 78, Austin, Texas.
- Hubbs, C.L., and G.P. Cooper. 1936. Minnows of Michigan. Cranbrook Institute of Science Bulletin 8.
- Hubbs, C.L., and K.F. Lagler. 1958. Fishes of the Great Lakes Region. University of Michigan Press, Ann Arbor, Michigan.
- Idaho Fish and Game. 1990. Fisheries Management Plan 1991-1995. Idaho Department of Fish and Game.
- Illinois Natural History Survey. 2004. Illinois Natural History Survey Fish Collection Database Search Results.
- Insider Viewpoint. 2001. *[Source did not provide reference]*.
- Jansen, C. 2003. Barrier to keep carp out of Big Muskego Lake. JS Online. Milwaukee Journal Sentinel. October 7, 2003.
- Jenkins, R.E., and N.M. Burkhead. 1994. Freshwater fishes of Virginia. American Fisheries Society, Bethesda, Maryland.
- Jia, Y., Y. Chen, S. Xie, and Y. Yang. 2008. Physiological advantages may contribute to successful invasion of the exotic *Cyprinus carpio* into the Xingyun Lake, China. Environmental Biology of Fishes. 81(4): 457-463.
- Johnson, M., and G.C. Becker. 1980. Annotated list of the fishes of Wisconsin. Papers of the Wisconsin Academy of Sciences, Arts and Letters 58: 265-300.
- Jones, M.J., and I.G. Stuart. 2006. Movement of common carp, *Cyprinus carpio*, in a regulated lowland Australian river: implications for management. Fisheries Management and Ecology. 13(4): 213-219.
- Jones, M.J., and I.G. Stuart. 2009. Lateral movement of common carp (*Cyprinus carpio* L.) in a large lowland river and floodplain. Ecology of Freshwater Fish. 18(1): 72-82.
- Jordan, D.S. 1882. Report on the fishes of Ohio. Report of the Geological Survey of Ohio 4(1): 735-1002.

- Jordan, D.S., and B.W. Evermann. 1902. Preliminary report on an investigation of the fishes and fisheries of the Hawaiian Islands. Pages 353-380 *in* Report of the Commissioner for the year ending June 30, 1901. U.S. Commission of Fish and Fisheries, Government Printing Office, Washington, District of Columbia.
- Jordan, D.S., and B.W. Evermann. 1905. The aquatic resources of the Hawaiian Islands. Part I. The shore fishes. Bulletin of the U.S. Fish Commission for 1903, 23: 1-574.
- Koehn, J.D. 2004. Carp (*Cyprinus carpio*) as a powerful invader in Australian waterways. *Freshwater Biology*. 49(7): 882-894.
- Koehn, J., A. Brumley, and P. Gehrke. 2000. Managing the Impacts of Carp. Bureau of Rural Sciences (Department of Agriculture, Fisheries and Forestry – Australia), Canberra.
- Koster, W.J. 1957. Guide to the Fishes of New Mexico. University of New Mexico Press, Albuquerque, New Mexico.
- Kottelat, M. 1997. European freshwater fishes. *Biologia* 52, Suppl. 5: 1-271.
- Kottelat, M. 2001. Fishes of Laos. WHT Publications Ltd., Colombo 5, Sri Lanka.
- Kuhne, E.R. 1939. A guide to the fishes of Tennessee and the mid-South. Tennessee Department of Conservation, Nashville, Tennessee.
- Laird, C.A., and L.M. Page. 1996. Non-native fishes inhabiting the streams and lakes of Illinois. *Illinois Natural History Survey Bulletin* 35(1): 1-51.
- Lamarra, V.A.J. 1975. Digestive activities of carp as a major contributor to the nutrient loading of lakes. *Proceedings of the International Association of Theoretical and Applied Limnology* 19: 2461–2468.
- Lampman, B.H. 1946. The coming of the pond fishes. Binfords and Mort, Portland, Oregon.
- La Rivers, I. 1962. Fishes and Fisheries of Nevada. Nevada State Print Office, Carson City, Nevada.
- Lee, D.S., A. Norden, C.R. Gilbert, and R. Franz. 1976. A list of the freshwater fishes of Maryland and Delaware. *Chesapeake Science* 17(3): 205-211.
- Lee, D.S., C.R. Gilbert, C.H. Hocutt, R.E. Jenkins, D.E. McAllister, and J.R. Stauffer, Jr. 1980 et seq. Atlas of North American freshwater fishes. North Carolina State Museum of Natural History, Raleigh, North Carolina.
- Linder, A.D. 1963. Idaho's alien fishes. *Tebiwa* 6(2): 12-15.

- Logan, D.J., E.L. Bibles, and D.F. Markle. 1996. Recent collections of continental exotic aquarium fishes in Oregon and thermal tolerance of Oriental weatherfish and pirapatinga. *California Fish and Game* 82(2): 66-80.
- Loyacano, H.A., Jr. 1975. A list of freshwater fishes of South Carolina. *Bulletin of the South Carolina Experimental Station* 580: 1-8.
- Machacek, H. 2007. World Records Freshwater Fishing. Available: www.fishing-worldrecords.com. (November 2007).
- Maciolek, J.A. 1984. Exotic fishes in Hawaii and other islands of Oceania. Pages 131-161 in W. R. Courtenay, Jr., and J. R. Stauffer, Jr., editors. *Distribution, biology, and management of exotic fishes*. The Johns Hopkins University Press, Baltimore, Maryland.
- Matern, S.A., P.B. Moyle, and L.C. Pierce. 2002. Native and alien fishes in a California estuarine marsh: twenty-one years of changing assemblages. *Transactions of the American Fisheries Society* 131: 797-816.
- McCarraher, D.B., and R.W. Gregory. 1970. Adaptability and status of introductions of Sacramento perch, *Archoplites interruptus*, in North America. *Transactions of the American Fisheries Society* 99(4): 700-707.
- McDonald, M. 1886. Report on distribution of fish and eggs by the U.S. Fish Commission for the season of 1885-'86. *Bulletin of the United States Fish Commission* 6(1886): 385-394.
- McDonald, M. 1893. Report of the Commissioner for 1889 to 1891. Part XVII. U.S. Commission of Fish and Fisheries, Washington, District of Columbia.
- Menhinick, E.F. 1991. *The Freshwater Fishes of North Carolina*. North Carolina Wildlife Resources Commission.
- Mettee, M.F., E. O'Neil, R.D. Suttkus, and J.M. Pierson. 1987. *Fishes of the Lower Tombigbee River System in Alabama and Mississippi*. Geological Survey of Alabama Bulletin 107.
- Mettee, M.F., P.E. O'Neil, and J.M. Pierson. 1996. *Fishes of Alabama and the Mobile Basin*. Oxmoor House, Inc. Birmingham, Alabama.
- Miller, R.R. 1952. Bait fishes of the lower Colorado River, from Lake Mead, Nevada, to Yuma, Arizona, with a key for identification. *California Fish and Game* 38: 7-42.
- Miller, R.R., and J.R. Alcorn. 1946. The introduced fishes of Nevada, with a history of their introduction. *Transactions of the American Fisheries Society* 73: 173-193.
- Miller, S.A., and T.A. Crowl. 2006. Effects of common carp (*Cyprinus carpio*) on macrophytes and invertebrate communities in a shallow lake. *Freshwater Biology* 51: 85-94.

- Miller, S.A., and F.D. Provenza. 2007. Mechanisms of resistance of freshwater macrophytes to herbivory by invasive juvenile common carp. *Freshwater Biology* (2007) 52: 39–49.
- Miller, R.R., and C.H. Lowe. 1967. Part 2. Fishes of Arizona. Pages 133-151 *in* C.H. Lowe, editor. *The Vertebrates of Arizona*. University of Arizona Press. Tucson, Arizona.
- Miller, R.J., and H.W. Robison. 1973. *The Fishes of Oklahoma*. Oklahoma State University Press, Stillwater, Oklahoma.
- Miller, A.I., and L.G. Beckman. 1996. First record of predation on white sturgeon eggs by sympatric fishes. *Transactions of the American Fisheries Society* 125: 338-340.
- Minckley, W.L. 1973. *Fishes of Arizona*. Arizona Fish and Game Department. Sims Printing Company, Inc., Phoenix, Arizona.
- Minnesota Sea Grant. 2004. Dumping of aquarium fish causing trouble in Duluth (or Something's fishing in Rock Pond). Available:
<http://www.seagrants.mn.edu/news/2004/may11.html>.
- Mississippi Museum of Natural Science. 2004. Mississippi Museum of Natural Science Nonindigenous Fish Records.
- Moore, H.H., and R.A. Bream. 1965. Distribution of fishes in U.S. streams tributary to Lake Superior. U.S. Fish and Wildlife Service Special Science Report 516.
- Morris, J., L. Morris, and L. Witt. 1974. *The Fishes of Nebraska*. Nebraska Game and Parks Commission Lincoln, Nebraska.
- Moyle, P.B. 1976. *Inland Fishes of California*. University of California Press, Berkeley, California.
- Moyle, P.B. 2002. *Inland Fishes of California*. Second Edition. University of California Press. Berkeley and Los Angeles, California.
- Moyle, J.B., and W.D. Clothier. 1959. Effects of management and winter oxygen levels on the fish population of a prairie lake. *Transactions of the American Fisheries Society* 88: 178-185.
- Moyle, P.B., F.W. Fisher, and H. Li. 1974. Mississippi silversides and logperch in the Sacramento-San Joaquin River system. *California Department of Fish and Game*. 60(2): 145-147.
- Moyle, P.B., and R.A. Daniels. 1982. Fishes of the Pit River System, McCloud River System, and Surprise Valley Region. University of California Publications, Zoology 115: 1-82.

- Mundy, B.C. 2005. Fishes of the Hawaiian Archipelago. Bishop Museum Bulletins in Zoology, Number 6.
- Myers, J. 2004. Drain a pond, save a stream. DuluthNewsTribune.com. May 10, 2004.
- National Institute of Water and Atmospheric Research (NIWA). 2009. Home › Our Science › Freshwater › Management Tools › NIWA Atlas of NZ Freshwater Fishes › Koi Carp Available: http://www.niwa.co.nz/our-science/freshwater/tools/fishatlas/species/koi_carp. (November 2009).
- Nelson, J. 1890. Descriptive catalogue of the vertebrates of New Jersey. Geological Survey of New Jersey 1890: 489-824.
- Nelson, J.S., and S.D. Gerking. 1968. Annotated key to the fishes of Indiana. Project 342-303-815. Department of Zoology, Indiana Aquatic Research Unit, Indiana State University, Bloomington, Indiana.
- New Mexico Game and Fish. 2000. Biota Information System of New Mexico BISON: Common carp, *Cyprinus carpio*. Available: http://www.fw.vt.edu/fishesex/nmex_main/species/010080.htm.
- New South Wales (NSW) Department of Primary Industries: Fishing and Aquaculture. 2005. Carp (*Cyprinus carpio*). Available: <http://www.dpi.nsw.gov.au/fisheries/pests-diseases/freshwater-pests/species/carp>. (November 2009).
- Nico, L. G. 2005. Changes in the fish fauna of the Kissimmee River Basin, peninsular Florida: nonnative additions. Pages 523-556 in J. N. Rinne, R. M. Hughes, and B. Calamusso, editors. Historical changes in large river fish assemblages of the Americas. American Fisheries Society Symposium 45, Bethesda, Maryland.
- Nico, L., E. Maynard, and P.J. Schofield. 2009. *Cyprinus carpio*. USGS Nonindigenous Aquatic Species Database, Gainesville, Florida. Available: <http://nas.er.usgs.gov/queries/FactSheet.asp?speciesID=4>. (November 2009).
- Nicol, S., J. Lieschke, J. Lyon, and J. Koehn. 2004. Observations on the distribution and abundance of carp and native fish, and their responses to a habitat restoration trial in the Murray River, Australia. New Zealand Journal of Marine and Freshwater Research 38: 541–551.
- Nunn, A., J.D. Bolland, J.P. Harvey, and I.G. Cowx. 2007. Establishment of self-sustaining populations of non-native fish species in the River Trent and Warwickshire Avon, UK, indicated by the presence of 0+ fish. Aquatic Invasions 2(3): 190-196.
- Owen, J.B., D.S. Elsen, and G.W. Russell. 1981. Distribution of Fishes in North and South Dakota Basins Affected by the Garrison Diversion Unit. Fisheries Research Unit, University of North Dakota, Grand Forks, North Dakota.

- Pearson, W.D., and L.A. Krumholz. 1984. Distribution and status of Ohio River fishes. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Pflieger, W.L. 1971. A distributional study of Missouri fishes. University of Kansas Publications, Museum of Natural History 20(3): 225-570.
- Pflieger, W.L. 1975. The Fishes of Missouri. Missouri Department of Conservation. Jefferson City, Missouri.
- Pflieger, W.L. 1997. The Fishes of Missouri. Missouri Department of Conservation, Jefferson City, Missouri.
- Phillips, G.L., W.D. Schmid, and J.C. Underhill. 1982. Fishes of the Minnesota Region. University of Minnesota Press, Minneapolis, Minnesota.
- Pinto, L., N. Chandrasena, J. Pera, P. Hawkins, D. Eccles, and R. Sim. 2005. Managing invasive carp (*Cyprinus carpio* L.) for habitat enhancement at Botany Wetlands, Australia. Aquatic Conservation: Marine and Freshwater Ecosystems 15: 447-462.
- Platania, S.P. 1991. Fishes of the Rio Chama and upper Rio Grande, New Mexico, with preliminary comments on their longitudinal distribution. Southwestern Naturalist 36(2): 186-193.
- Power, G.J., and F. Ryckman. 1998. North Dakota Fisheries Investigations: status of North Dakota's fishes. Available: <http://www.npwrc.usgs.gov/resource/fish/fshries/index.htm>.
- Powers, S.L., and P.A. Ceas. 2000. Ichthyofauna and biogeography of Russell Fork (Big Sandy River - Ohio River). Southeastern Fishes Council Proceedings 41: 1-12.
- Raasch, M.S., and V.L. Altemus, Sr. 1991. Delaware's freshwater and brackish water fishes - a popular account. Delaware State College for the Study of Del-Mar-Va Habitats and the Society of Natural History of Delaware
- Rasmussen, J.L. 1998. Aquatic nuisance species of the Mississippi River basin. 60th Midwest Fish and Wildlife Conference, Aquatic Nuisance Species Symposium, Dec. 7, 1998, Cincinnati, Ohio.
- Ravenel, W.C. 1896. Report on the propagation and distribution of food-fishes. Pages 6-72 in Report of the Commissioner for the year ending June 30, 1895, Part XXI. U.S. Commission of Fish and Fisheries, Washington, District of Columbia.
- Ravenel, W.C. 1898. Report on the propagation and distribution of food-fishes. Pages 11-92 in Report of the Commissioner for the year ending June 30, 1896, Part XXII. U.S. Commission of Fish and Fisheries, Washington, District of Columbia.

- Red River Authority of Texas. 2001. Red and Canadian Basins Fish Inventory: Grayson County. Red River Authority of Texas.
- Redding, J.D. 1884. Character of the carp introduced by Capt. Henry Robinson about 1830. Bulletin of the U.S. Fish Commission 4: 266-267.
- Richardson, M.J., F.G. Whoriskey, and L.H. Roy. 1995. Turbidity generation and biological impacts of an exotic fish *Carassius auratus*, introduced into shallow seasonally anoxic ponds. Journal of Fish Biology 47: 576-585.
- Riede, K. 2004. Global register of migratory species - from global to regional scales. Final Report of the R&D-Projekt 808 05 081. Federal Agency for Nature Conservation, Bonn, Germany.
- Roberts, J., A. Chick, L. Oswald, and P. Thompson. 1995. Effect of Carp, *Cyprinus carpio* L., an Exotic Benthivorous Fish, on Aquatic Plants and Water Quality in Experimental Ponds. Marine: Freshwater Research 46: 1171-80.
- Robison, H.W., and T.M. Buchanan. 1988. Fishes of Arkansas. University of Arkansas Press. Fayetteville, Arkansas.
- Ross, S.T., and W.M. Brennenman. 1991. Distribution of freshwater fishes in Mississippi. Manuscript. Mississippi Department of Wildlife, Fisheries and Parks, Jackson, Mississippi.
- Rule, R. 1885. Southern Region. Pages 10-15 in J. H. Taggart, editor. Annual Report of the Arizona Fish Commission, 1883-1884, to Frederick A. Tritle, Governor of the Territory of Arizona.
- Ryon, M.G., and J.M. Loar. 1988. Checklist of fishes on the Department of Energy Oak Ridge Reservation. Journal of the Tennessee Academy of Sciences 63: 97-102.
- Saikia, S.K., and D.N. Das. 2009. Feeding ecology of common carp (*Cyprinus carpio* L.) in a rice-fish culture system of the Apatani plateau (Arunachal Pradesh, India). Aquatic Ecology 43(2): 559-568.
- Scarola, J.F. 1973. Freshwater Fishes of New Hampshire. New Hampshire Fish and Game Department, Division of Inland and Marine Fisheries.
- Scheffer, M. 1998. Ecology of shallow lakes. Chapman and Hall, London.
- Schramm, H.L., Jr., and M.C. Basler. 2004. Evaluation of capture methods and distribution of black carp in Mississippi. Mississippi State University, Starkville, Mississippi.
- Schwartz, F. 1963. The fresh-water minnows of Maryland. Maryland Conservationist 40(2): 19-29.

- Scoppettone, G.G., P.H. Rissler, M.B. Nielsen, and J.E. Harvey. 1998. The status of *Moapa coriacea* and *Gila seminuda* and status information on other fishes of the Muddy River, Clark County, Nevada. *Southwestern Naturalist* 43(2): 115-122.
- Scott, W.B., and E.J. Crossman. 1973. *Freshwater Fishes of Canada*. Fisheries Research Board of Canada, Ottawa. Bulletin 184.
- Shafland, P.L. 1996. Exotic fishes of Florida – 1994. *Reviews in Fisheries Science* 4(2): 101-122.
- Shebley, W.H. 1917. History of the introduction of food and game fishes into the waters of California. *California Fish and Game* 3(1): 3-10.
- Shields, J.T. 1958a. Experimental control of carp reproduction through water drawdowns in Fort Randall Reservoir, South Dakota. *Transactions of the American Fisheries Society* 87: 23-33.
- Shields, J.T. 1958b. Fish management problems of large impoundments on the Missouri River. *Transactions of the American Fisheries Society* 87: 356-362.
- Shin-ichiro et al. 1009. *[Source did not provide references]*.
- Sigler, W.F., and R.R. Miller. 1963. *Fishes of Utah*. Utah Department of Fish and Game, Salt Lake City, Utah.
- Sigler, W.F., and J.W. Sigler. 1996. *Fishes of Utah*. University of Utah Press, Salt Lake City, Utah.
- Simpson, J., and R. Wallace. 1978. *Fishes of Idaho*. University of Idaho Press, Moscow, Idaho.
- Smiley, C.W. 1886. Some results of carp culture in the United States. Pages 657-890 in Report of the Commissioner of Fish and Fisheries for 1884, Part XII. U.S. Commission of Fish and Fisheries, Washington, District of Columbia.
- Smith, P.W. 1979. *The Fishes of Illinois*. University of Illinois Press, Urbana, Illinois.
- Smith, J.J. 1982. Fishes of the Pajaro River System. Pages 83-170 in P. B. Moyle, J. J. Smith, R. A. Daniels, T. L. Price, and D. M. Baltz, editors. *Distribution and ecology of stream fishes of the Sacramento-San Joaquin drainage system, California*. University of California Press, Berkeley, California.
- Smith, H.M. 1896. A review of the history and results of the attempts to acclimatize fish and other water animals in the Pacific states. *Bulletin of the U.S. Fish Commission for 1895*, 40: 379-472.

- Smith-Vaniz, W.F. 1968. Freshwater fishes of Alabama. Auburn University Agricultural Experiment Station, Auburn, Alabama.
- Sommer, T., B. Harrell, M. Nobriga, R. Brown, P. Moyle, W. Kimmerer, and L. Schemel. 2001. California's Yolo Bypass: Evidence that flood control can be compatible with fisheries, wetlands, wildlife, and agriculture. *Fisheries* 26 (8): 6-16.
- South Australia Government: Department for Environment and Heritage. Undated. Supporting Native Fish: Backyards for Wildlife Fact Sheet.
- Spencer, S.L., W.E. Swingle, and T. Scott, Jr. 1964. Commercial fishing in the Mobile Delta, Alabama, during the period of July 1, 1963 to June 30, 1964. Alabama Department of Conservation 1:17.
- Spillman, C.J. 1961. Faune de France: Poissons d'eau douce. Fédération Française des Sociétés Naturelles, Tome 65. Paris.
- Starnes, W.C., J. Odenkirk, and M.J. Ashton. 2011. Update and analysis of fish occurrences in the lower Potomac River drainage in the vicinity of Plummerville Island, Maryland—Contribution XXXI to the natural history of Plummerville Island, Maryland. *Proceedings of the Biological Society of Washington* 124(4): 280-309.
- Stauffer, J.R., Jr., J.M. Boltz, and L.R. White. 1995. The Fishes of West Virginia. West Virginia Department of Natural Resources. Academy of Natural Sciences of Philadelphia, Philadelphia, Pennsylvania.
- Stiles, E.W. 1978. Vertebrates of New Jersey. Edmund W. Stiles, Somerset, New Jersey.
- Stone, M.D. 1995. Fish stocking programs in Wyoming: a balanced perspective. Pages 47-51 in H.L. Schramm, Jr., and R.G. Piper, editors. Uses and effects of cultured fishes in aquatic ecosystems. American Fisheries Society Symposium 15.
- Sublette, J.E., M.D. Hatch, and M. Sublette. 1990. The Fishes of New Mexico. New Mexico Department of Game and Fish, University of New Mexico Press, Albuquerque, New Mexico.
- Suzuki, R., and M. Yamaguchi. 1980. Meristic and morphometric characters of five races of *Cyprinus carpio*. *Japanese Journal of Ichthyology* 27(3): 199-206.
- Suzuki, R., and M. Yamaguchi. 1984. Meristic and morphometric characters of interracial hybrids of the common carp *Cyprinus carpio*. *Bulletin of the National Research Institute of Aquaculture* 6: 1-9.
- Sweeney, Z.T. 1902. Biennial Report of the Commissioner of Fisheries and Game for Indiana. Indianapolis, Indiana.

- Swift, C., R.W. Yerger, and P.R. Parrish. 1977. Distribution and natural history of the fresh and brackish water fishes of the Ochlockonee River, Florida and Georgia. Bulletin of Tall Timbers Research Station 20.
- Taggart, J.H. 1885. Annual Report of the Arizona Fish Commission, 1883-1884, to Frederick A. Tittle, Governor of the Territory of Arizona.
- Tanner, V.M. 1936. A study of the fishes of Utah. Utah Academy of Sciences, Arts and Letters 13: 155-183.
- Taylor, J.N., W.R. Courtenay, Jr., and J.A. McCann. 1984. Known impact of exotic fishes in the continental United States. Pages 322-373 in W.R. Courtenay, Jr., and J.R. Stauffer, editors. Distribution, biology, and management of exotic fish. Johns Hopkins Press, Baltimore, Maryland.
- Texas Parks and Wildlife Department. 1993. State Record Listing. May 21, 1993: 19-21.
- Texas Parks and Wildlife Department. 1994. *[Source did not provide reference]*.
- Texas Parks and Wildlife Department. 2001. Fish Records: Water Body - All Tackle. Texas Parks and Wildlife Department. April 24, 2001.
- Texas System of Natural Laboratories, Inc. 1996. Fish of Colorado River Basin, Texas. Texas System of Natural Laboratories, Inc.
- Texas System of Natural Laboratories, Inc., and USGS. 1994. Fish collected in streams of the South-Central Texas study unit, Texas in Freshwater and marine fishes of Texas and the northwestern Gulf of Mexico: taxonomic and distributional inventory with bibliography. Texas System of Natural Laboratories, Inc.
- Tilmant, J.T. 1999. Management of nonindigenous aquatic fish in the U.S. National Park System. National Park Service.
- Trautman, M.B. 1981. The Fishes of Ohio. Ohio State University Press, Columbus, Ohio.
- Truitt, R.V., B.A. Bean, and H.W. Fowler. 1929. The Fishes of Maryland. Maryland Conservation Department, Conservation Bulletin 3, Annapolis, Maryland.
- Tyus, H.M., and J.F. Saunders. 2000. Non-native Fish Control and Endangered Fish Recovery: Lessons from the Colorado River. Fisheries 25(9): 17-24.
- Underhill, J.C. 1959. Fishes of the Vermillion River, South Dakota. Proceedings of the South Dakota Academy of Science 38: 96-102.
- University of Michigan Museum of Zoology. 2004. Michigan Fish Atlas Maps. Available: http://www.michigan.gov/cgi/0,1607,7-158-12540_13085_13553-30538--,00.html.

- U.S. Fish and Wildlife Service (USFWS). 2005. National Wildlife Refuge System Invasive Species. Available: <http://www.nwrinvasives.com/index.asp>. (2006).
- Vanicek, C.D., R.H. Kramer, and D.R. Franklin. 1970. Distribution of Green River fishes in Utah and Colorado following closure of Flaming Gorge Dam. *The Southwestern Naturalist* 14(3): 297-315.
- Vinyard, G.L. 2001. Fish Species Recorded from Nevada. Biological Resources Research Center 5. Available: <http://www.brrc.unr.edu/data/animal/vertebrates/fishlist.htm>.
- Walters, D.M. 1997. The distribution, status, and ecology of the fishes of the Conasauga River system. Master's Thesis, University of Georgia, Athens, Georgia.
- Webster, D.A. 1941. The life histories of some Connecticut fishes. Pages 122-227 *in* State Board of Fisheries and Game. A fishery survey of important Connecticut lakes. Connecticut Geological and Natural History Survey 63.
- Werner, R.G. 1980. Freshwater fishes of New York state: a field guide. Syracuse University Press, Syracuse, New York.
- Whitworth, W.R. 1996. Freshwater Fishes of Connecticut. State Geological and Natural History Survey of Connecticut, Bulletin 114.
- Wiltzius, W.J. 1981. Compendium of introduction date and state and federal annual stocking of various fishes in Colorado, 1972-1978. Colorado Division of Wildlife, Fort Collins, Colorado. Unpublished report.
- Woodling, J. 1985. Colorado's little fish: a guide to the minnows and other lesser known fishes in the state of Colorado. Colorado Division of Wildlife, Denver, Colorado.
- Worth, S.G. 1895. Report on the propagation and distribution of food-fishes. Pages 78-138 *in* Report of the Commissioner for the year ending June 30, 1893, Part XIX. U.S. Commission of Fish and Fisheries, Washington, District of Columbia.
- Wydoski, R.S., and R.R. Whitney. 1979. Inland Fishes of Washington. University of Washington Press, Seattle, Washington.
- Young, B.A., T.L. Welker, M.L. Wildhaber, C.R. Berry, and D. Scarnecchia, editors. 1997. Population structure and habitat use of benthic fishes along the Missouri and lower Yellowstone rivers. Annual Report of Missouri River Benthic Fish Study PD-95-5832. U.S. Army Corps of Engineers and the U.S. Bureau of Reclamation.
- Zambrano, L., M.R. Perrow, C. Macias-Garcia, and V. Aguirre-Hidalgo. 1999. Impact of introduced carp (*Cyprinus carpio*) in subtropical shallow ponds in Central Mexico. *Journal of Aquatic Ecosystem Stress and Recovery* 6: 281-288.

Zuckerman, L.D., and R.J. Behnke. 1986. Introduced fishes in the San Luis Valley, Colorado. Pages 435-453 *in* R.H. Stroud, editor. Fish Culture In Fisheries Management. Proceedings of a symposium on the role of fish culture in fisheries management at Ozark, MO. American Fisheries Society, Bethesda, Maryland.